arXiv:1403.2195v1 [math.CA] 10 Mar 2014

A generalized integral transform and an alternative technique for solving linear ordinary differential equations

Nese Dernek[1] * and Fatih Aylikci[2]

[1]Department of Mathematics, Marmara University, Istanbul, Turkey
[2]Department of Mathematical Engineering, Yildiz Technical University, Istanbul, Turkey

March 11, 2014

Abstract

In the present paper authors introduce the $\mathcal{L}_n$-integral transform and the $\mathcal{L}_n^{-1}$ inverse integral transform for $n = 2^k$, $k \in \mathbb{N}$, as a generalization of the classical Laplace transform and the $\mathcal{L}^{-1}$ inverse Laplace transform, respectively. Applicability of this transforms in solving linear ordinary differential equations is analyzed. Some illustrative examples are also given.

Keywords: The Laplace transform, The $\mathcal{L}_2$-transform,The $\mathcal{L}_n$-transform, The $\mathcal{L}_n^{-1}$-transform and Linear ordinary differential equations.

1. Introduction, definitions and preliminaries

The following Laplace-type the $\mathcal{L}_2$ transform

$$\mathcal{L}_2\{f(x);y\} = \int_0^\infty x \exp(-x^2y^2) f(x)dx \tag{1.1}$$

was introduced by Yurekli and Sadek [4]. After then Aghili, Ansari and Sedghi [1] derived a complex inversion formula as follows

$$\mathcal{L}_2^{-1}\{\mathcal{L}_2\{f(x);y\}\} = \frac{1}{2\pi i}\int_{c-i\infty}^{c+i\infty} 2\mathcal{L}_2\{f(x);\sqrt{y}\}\exp(yx^2)dy \tag{1.2}$$

*Correspondence: ndernek@marmara.edu.tr
2010 AMS Mathematics Subject Classification: 44A10, 44A15, 44A20, 34A30

where $\mathcal{L}_2\{f(x);\sqrt{y}\}$ has a finite number of singularities in the left half plane $Re\ y \leqslant c$.

In this article, we introduce a new generalized integral transform as follows

$$\mathcal{L}_n\{f(x);y\} = \int_0^\infty x^{n-1}\exp(-x^n y^n) f(x)dx \tag{1.3}$$

where $n = 2^k,\ \ k \in \mathbb{N}$.

The $\mathcal{L}_n$-transform is related to the Laplace transform by means

$$\mathcal{L}_n\{f(x);y\} = \frac{1}{n}\mathcal{L}\{f(x^{\frac{1}{n}});y^n\}, \tag{1.4}$$

where the Laplace transform is defined by

$$\mathcal{L}\{f(x);y\} \int_0^\infty \exp(-xy) f(x)dx. \tag{1.5}$$

First we shall give several examples of the $\mathcal{L}_n$-transforms of some elementary and special functions.

Example 1.1. We show that

$$\mathcal{L}_n\{1;y\} = \int_0^\infty x^{n-1}\exp(-y^n x^n)dx = \frac{1}{ny^n}. \tag{1.6}$$

Example 1.2. We show that

$$\mathcal{L}_n\{x^k;y\} = \frac{1}{ny^{n+k}}\Gamma(\frac{k}{n}+1) \tag{1.7}$$

where $k, n \in \mathbb{N}$ and $k > -n$. Applying the definition of the $\mathcal{L}_n$-transform, we have

$$\mathcal{L}_n\{x^k;y\} = \int_0^\infty x^{k+n-1}\exp(-y^n x^n)dx, \tag{1.8}$$

where $k \in \mathbb{N}$.

The integral on the right hand side may be evaluated by changing the variable of the integration from x to u where $x^n y^n = u$, and using Gamma function's relation in (1.8), we obtain

$$\mathcal{L}_n\{x^k;y\} = \frac{1}{ny^{n+k}}\int_0^\infty u^{\frac{k}{n}+1-1}\exp(-u)du = \frac{1}{ny^{n+k}}\Gamma(\frac{k}{n}+1). \tag{1.9}$$

Example 1.3. We show that

$$\mathcal{L}_n\{\cos(ax^n);y\} = \frac{y^n}{n(y^{2n}+a^2)}. \tag{1.10}$$

Using the definition of the $\mathcal{L}_n$-transform and calculating the Taylor expansion of the $\cos$ function in (1.10) we get

$$\mathcal{L}_n\{\cos(ax^n); y\} = \sum_{m=0}^{\infty}(-1)^m \frac{a^{2m}}{(2m)!}\mathcal{L}_n\{x^{2mn}; y\}, \tag{1.11}$$

where from (1.7) we have the following relation

$$\mathcal{L}_n\{x^{2mn}; y\} = \frac{2m+1}{ny^{n+2mn}} \tag{1.12}$$

and then we obtain (1.10)

$$\mathcal{L}_n\{\cos(ax^n); y\} = \frac{1}{ny^n}\sum_{m=0}^{\infty}(-1)^m \frac{a^{2m}}{y^{2mn}} = \frac{y^n}{n(y^{2n}+a^2)}. \tag{1.13}$$

Example 1.4. We show that

$$\mathcal{L}_n\{\sin(ax^n); y\} = \frac{a}{n(y^{2n}+a^2)}. \tag{1.14}$$

Using the linearity of the $\mathcal{L}_n$-transform and calculating the Taylor expansion of the $\sin$ function in (1.14) we get

$$\mathcal{L}_n\{\sin(ax^n); y\} = \sum_{m=0}^{\infty}(-1)^m \frac{a^{2m+1}}{(2m+1)!}\mathcal{L}_n\{x^{(2m+1)n}; y\}. \tag{1.15}$$

Using the following relation in (1.7) of Example 1.2,

$$\mathcal{L}_n\{x^{(2m+1)n}; y\} = \frac{\Gamma(2m+2)}{ny^{2m+2n}}, \tag{1.16}$$

we have

$$\mathcal{L}_n\{\sin(ax^n); y\} = \frac{a}{ny^{2n}}\sum_{m=0}^{\infty}\frac{(-a^2)^m}{y^{2mn}} = \frac{a}{n(y^{2n}+a^2)}. \tag{1.17}$$

Example 1.5. We show that

$$\mathcal{L}_n\{\exp(-a^n x^n); y\} = \frac{1}{n(y^n+a^n)} \tag{1.18}$$

where $0 < Re(a) < Re(y)$.

Using the definition of the $\mathcal{L}_n$-transform and calculating the Taylor expansion of the exponential function we have

$$\mathcal{L}_n\{\exp(-a^n x^n); y\} = \sum_{m=0}^{\infty}(-1)^m \frac{a^{mn}}{m!}\mathcal{L}_n\{x^{mn}; y\}. \tag{1.19}$$

Using the value

$$\mathcal{L}_n\{x^{mn}; y\} = \frac{\Gamma(m+1)}{ny^{n+mn}} \tag{1.20}$$

on the right hand side of (1.19) we get

$$\mathcal{L}_n\{\exp(-a^n x^n); y\} = \frac{1}{ny^n}\sum_{m=0}^{\infty}(-1)^m \frac{a^{mn}}{y^{mn}} = \frac{1}{n(y^n+a^n)}. \tag{1.21}$$

Example 1.6. We show that

$$\mathcal{L}_n\{J_0(2a^{\frac{n}{2}}x^{\frac{n}{2}});y\} = \frac{1}{ny^n}\exp(-\frac{a^n}{y^n}) \tag{1.22}$$

where the function J_0 is the Bessel function of the first kind of order zero.

Using the following Taylor expansion of the function $J_0(x)$,

$$J_0(x) = \sum_{m=0}^{\infty}\frac{(-1)^m}{(m!)^2}\Big(\frac{x}{2}\Big)^{2m}, \tag{1.23}$$

we obtain

$$\mathcal{L}_n\{J_0(2a^{\frac{n}{2}}x^{\frac{n}{2}});y\} = \sum_{m=0}^{\infty}\frac{(-1)^m a^{mn}}{m!\Gamma(m+1)}\mathcal{L}_n\{x^{mn};y\}. \tag{1.24}$$

We know the $\mathcal{L}_n$ transform of $f(x) = x^{mn}$ as

$$\mathcal{L}_n\{x^{mn};y\} = \frac{\Gamma(m+1)}{ny^{n+mn}}. \tag{1.25}$$

Substituting the relation 1.25 into (1.24) we obtain

$$\mathcal{L}_n\{J_0(2a^{\frac{n}{2}}x^{\frac{n}{2}});y\} = \frac{1}{ny^n}\sum_{m=0}^{\infty}\frac{(-\frac{a^n}{y^n})^m}{m!} = \frac{1}{ny^n}\exp(-\frac{a^n}{y^n}). \tag{1.26}$$

Example 1.7. We show that

$$\mathcal{L}_n\{x^{\frac{nv}{2}}J_v(2a^{\frac{n}{2}}x^{\frac{n}{2}});y\} = \frac{1}{n}a^{\frac{nv}{2}}y^{-n(v+1)}\exp(-\frac{a^n}{y^n}) \tag{1.27}$$

where $Re(a) > 0, Rev > -1$.

Using the following Taylor expansion of $J_v(x)$, which is the Bessel function of the first kind of order v,

$$J_v(x) = \sum_{m=0}^{\infty}\frac{(-1)^m}{m!\Gamma(m+v+1)}\Big(\frac{x}{2}\Big)^{2m+v}, \tag{1.28}$$

we obtain

$$\mathcal{L}_n\{x^{\frac{nv}{2}}J_v(2a^{\frac{n}{2}}x^{\frac{n}{2}});y\} = \sum_{m=0}^{\infty}\frac{(-1)^m a^{nm+\frac{nv}{2}}}{m!\Gamma(m+v+1)}\mathcal{L}_n\{x^{mn+nv};y\}. \tag{1.29}$$

We can calculate the $\mathcal{L}_n$ transform of $f(x) = x^{mn+nv}$ function as follows

$$\mathcal{L}_n\{x^{mn+nv};s\} = \frac{\Gamma(m+v+1)}{ny^{n+mn+nv}}. \tag{1.30}$$

Substituting the relation 1.30 into equation (1.29) we obtain the assertion (1.27) of Example 1.7,

$$\mathcal{L}_n\{x^{\frac{nv}{2}}J_v(2a^{\frac{n}{2}}x^{\frac{n}{2}});y\} = \frac{a^{\frac{nv}{2}}}{ny^{nv+n}}\sum_{m=0}^{\infty}\frac{(-\frac{a^n}{y^n})^m}{m!} = \frac{1}{n}a^{\frac{nv}{2}}y^{-n(v+1)}\exp(-\frac{a^n}{y^n}). \tag{1.31}$$

Example 1.8. We show that

$$\mathcal{L}_n\{erfc(\frac{1}{2}\, a^{\frac{n}{2}}x^{-\frac{n}{2}}); y\} = \frac{1}{n}y^{-n}\exp(-a^{\frac{n}{2}}y^{\frac{n}{2}}) \tag{1.32}$$

where $Re(a) > 0$.

Using the definition of the complementary error function $erfc(x)$,

$$erfc(x) = \frac{2}{\sqrt{\pi}}\int_x^\infty \exp(-u^2)du, \tag{1.33}$$

we get

$$\mathcal{L}_n\{erfc(\frac{1}{2}\, a^{\frac{n}{2}}x^{-\frac{n}{2}}); y\} = \frac{2}{\sqrt{\pi}}\int_0^\infty x^{n-1}\exp(-y^n x^n)\int_{\frac{a^{n/2}}{2x^{n/2}}}^\infty \exp(-u^2)dudx. \tag{1.34}$$

Changing the order of integration, we obtain

$$\mathcal{L}_n\{erfc(\frac{1}{2}\, a^{\frac{n}{2}}x^{-\frac{n}{2}}); y\} = \frac{2}{\sqrt{\pi}}\int_0^\infty \exp(-u^2)\int_{\frac{a}{2^{2/n}u^{2/n}}}^\infty x^{n-1}\exp(-y^n x^n)dxdu \tag{1.35}$$

and using the relation $\frac{d}{dx}(\exp(-y^n x^n)) = -ny^n x^{n-1}\exp(-y^n x^n)$ we have

$$\mathcal{L}_n\{erfc(\frac{1}{2}\, a^{\frac{n}{2}}x^{-\frac{n}{2}}); y\} = \frac{2}{\sqrt{\pi}ny^n}\int_0^\infty \exp(-u^2)\exp(-\frac{y^n a^n}{4u^2})du. \tag{1.36}$$

Changing the variable from u to x according to the transformation $u = \frac{a^{n/2}}{2x^{n/2}}$ we find that

$$\mathcal{L}_n\{erfc(\frac{1}{2}\, a^{\frac{n}{2}}x^{-\frac{n}{2}}); y\} = \frac{a^{\frac{n}{2}}}{2\sqrt{\pi}y^n}\mathcal{L}_n\{x^{-\frac{3n}{2}}\exp(-\frac{a^n}{4x^n}); y\}. \tag{1.37}$$

Using the Taylor expansion of exponential function and the $\mathcal{L}_n$-transform of $f(x) = x^{-mn-\frac{3n}{2}}$ we obtain

$$\frac{a^{\frac{n}{2}}}{2\sqrt{\pi}y^n}\mathcal{L}_n\{x^{-\frac{3n}{2}}\exp(-\frac{a^n}{4x^n}); y\} = \frac{a^{\frac{n}{2}}}{2\sqrt{\pi}y^n}\sum_{m=0}^\infty \frac{(-1)^m a^{mn}}{m!4^m}\mathcal{L}_n\{x^{-mn-\frac{3n}{2}}; y\}$$

$$= \frac{a^{\frac{n}{2}}}{2n\sqrt{\pi}y^n}\sum_{m=0}^\infty \frac{(-1)^m a^{mn}}{m!4^m}\frac{\Gamma(-m-\frac{3}{2}+1)}{y^{-mn-\frac{n}{2}}}. \tag{1.38}$$

From the following Euler's reflection formula,

$$\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin(\pi z)}, \tag{1.39}$$

we get

$$\frac{a^{\frac{n}{2}}}{2n\sqrt{\pi}y^n}\sum_{m=0}^{\infty}\frac{(-1)^m a^{mn}}{m!4^m}\frac{\Gamma(-m-\frac{3}{2}+1)}{y^{-mn-\frac{n}{2}}}=\frac{a^{\frac{n}{2}}\pi}{2n\sqrt{\pi}y^n}\sum_{m=0}^{\infty}\frac{(-1)^{2m+1}a^{mn}y^{mn+\frac{n}{2}}}{\Gamma(m+1)\Gamma(m+1+\frac{1}{2})4^m} \tag{1.40}$$

and using the following duplication formula for Gamma function

$$\Gamma(z)\Gamma(z+\frac{1}{2})=2^{1-2z}\sqrt{\pi}\,\Gamma(2z) \tag{1.41}$$

we obtain

$$\frac{a^{\frac{n}{2}}\pi}{2n\sqrt{\pi}y^n}\sum_{m=0}^{\infty}\frac{(-1)^{2m+1}a^{mn}y^{mn+\frac{n}{2}}}{\Gamma(m+1)\Gamma(m+1+\frac{1}{2})4^m}=\frac{1}{n}y^{-n}\exp(-a^{\frac{n}{2}}y^{\frac{n}{2}}). \tag{1.42}$$

Thus the assertion (1.32) follows from (1.42).

Example 1.9. We show that

$$\mathcal{L}_n\{erf(a^{\frac{n}{2}}x^{\frac{n}{2}});y\}=\frac{a^{\frac{n}{2}}}{n}y^{-n}(y^n+a^n)^{-\frac{1}{2}} \tag{1.43}$$

where $-Re(a)<y, Re(y)>0$.

Using the definition of $\mathcal{L}_n$-transform and the error function we have

$$\mathcal{L}_n\{erf(a^{\frac{n}{2}}x^{\frac{n}{2}});y\}=\frac{2}{\sqrt{\pi}}\int_0^{\infty}x^{n-1}\exp(-y^nx^n)\int_0^{a^{n/2}x^{n/2}}\exp(-u^2)dudx. \tag{1.44}$$

Changing the order of integration and evaluating the inner integral we get

$$\mathcal{L}_n\{erf(a^{\frac{n}{2}}x^{\frac{n}{2}});y\}=\frac{2}{\sqrt{\pi}}\int_0^{\infty}\exp(-u^2)\int_{\frac{u^{2/n}}{a}}^{\infty}x^{n-1}\exp(-y^nx^n)dxdu$$

$$=\frac{2}{\sqrt{\pi}ny^n}\int_0^{\infty}\exp(-u^2(1+\frac{y^n}{a^n}))du. \tag{1.45}$$

Changing the variable from u to x according to transformation $u\sqrt{1+\frac{y^n}{a^n}}=x,$ we obtain the assertion (1.43),

$$\frac{2}{\sqrt{\pi}ny^n}\int_0^{\infty}\exp(-u^2(1+\frac{y^n}{a^n}))du=\frac{a^{\frac{n}{2}}}{n}y^{-n}(y^n+a^n)^{-\frac{1}{2}}. \tag{1.46}$$

Example 1.10. We show that

$$\mathcal{L}_n\{\exp(-ax^{2n});y\}=\frac{\sqrt{\pi}}{2n\sqrt{a}}\,\exp(\frac{y^{2n}}{4a})erfc(\frac{y^{2n}}{2\sqrt{a}}) \tag{1.47}$$

provided that $Re\ a > 0$.

Using the definition of the $\mathcal{L}_n$-transform we get

$$\mathcal{L}_n\{\exp(-ax^{2n}); y\} = \int_0^\infty x^{n-1} \exp(-y^n x^n - ax^{2n}) dx. \tag{1.48}$$

Writing on the right hand side of (1.48)

$$-y^n x^n - ax^{2n} = -a(x^n + \frac{y^n}{2a})^2 + \frac{y^{2n}}{4a} \tag{1.49}$$

and changing the variable

$$a^{1/2}(x^n + \frac{y^n}{2a}) = u, \tag{1.50}$$

using the definition of the complementary error function as follows, we deduce the assertion (1.48),

$$\mathcal{L}_n\{\exp(-ax^{2n}); y\} = \frac{\sqrt{\pi}}{2n\sqrt{a}}\ \exp(\frac{y^{2n}}{4a}) erfc(\frac{y^n}{2\sqrt{a}}). \tag{1.51}$$

Corollary 1.1. From the definition of the $\mathcal{L}_n$-transform the following identity hold true:

$$\mathcal{L}_n\{\exp(-ax^n) f(x); y\} = \mathcal{L}_n\{f(x); (y^n + a)^{\frac{1}{n}}\} \tag{1.52}$$

where $Re\ a > 0$.

We now introduce a new derivative operator for the $\mathcal{L}_n$-transform and apply the operator to solve following ordinary differential equations:

$$xz'' - (2v + n - 3)z' + x^{n-1}z = 0\ \ ,\ \ n = 2^k, k \in \mathbb{N}, v > n, v = 2^m + 1, m \in \mathbb{N} \tag{1.53}$$

$$xz'' - (n^2 - 1)z' + x^{n-1}z = 0\ \ ,\ \ n = 2^k,\ k = 0, 1, 2, ... \tag{1.54}$$

2. Some properties of the $\mathcal{L}_n$-transform

In this section we will give some properties of the $\mathcal{L}_n$-transform that will be used to solve the ordinary differential equations (1.53)-(1.54) given above.

Firstly, we introduce a differential operator $\overline{\delta}$ (see [7, 8]) that we call the $\overline{\delta}$- derivative and define as

$$\overline{\delta}_x = \frac{1}{x^{n-1}} \frac{d}{dx},\ \ n = 2^k,\ k \in \mathbb{N} \tag{2.1}$$

we note that

$$\overline{\delta}_x^2 = \overline{\delta}_x \overline{\delta}_x = \frac{1}{x^{n-1}} \frac{d}{dx} \Big(\frac{1}{x^{n-1}} \frac{d}{dx} \Big) = \frac{1}{x^{2n-2}} \frac{d^2}{dx^2} - \frac{(n-1)}{x^{2n-1}} \frac{d}{dx}. \tag{2.2}$$

The $\overline{\delta}$ derivative operator can be successively applied in a similar fashion for any positive integer power.

Here we will derive a relation between the $\mathcal{L}_n$-transform of the $\overline{\delta}$-derivative of a function and the $\mathcal{L}_n$-transform of the function itself.

Suppose that $f(x)$ is a continuous function with a piecewise continuous derivative $f'(x)$ on the interval $[0,\infty)$. Also suppose that f and f' are of exponential order $\exp(\alpha^n x^n)$ as $x \to \infty$ where α is a constant. By using the definitions of $\mathcal{L}_n$-transform and the $\overline{\delta}$ derivative and integration by parts, we obtain

$$\mathcal{L}_n\{\overline{\delta}_x f(x); y\} = \int_0^\infty \exp(-y^n x^n) f'(x) dx, \tag{2.3}$$

$$\int_0^\infty \exp(-y^n x^n) f'(x) dx = \lim_{b\to\infty} f(x)\exp(-y^n x^n)|_0^b + ny^n \int_0^\infty x^{n-1}\exp(-y^n x^n) f(x) dx. \tag{2.4}$$

Since f is of exponential order $\exp(\alpha^n x^n)$ as $x \to \infty$, it follows that

$$\lim_{x\to\infty} \exp(-y^n x^n) f(x) = 0 \tag{2.5}$$

and consequently,

$$\mathcal{L}_n\{\overline{\delta}_x f(x); y\} = ny^n \mathcal{L}_n\{f(x); y\} - f(0^+). \tag{2.6}$$

Similarly, if f and f' are continuous functions with a piecewise continuous derivative f'' on the interval $[0,\infty)$, and if all three functions are of exponential order $\exp(\alpha^n x^n)$ as $x \to \infty$ we can use (2.6) to obtain

$$\mathcal{L}_n\{\overline{\delta}_x^2 f(x); y\} = n^2 y^{2n} \mathcal{L}_n\{f(x); y\} - ny^n f(0^+) - \overline{\delta}_x f(0^+). \tag{2.7}$$

Using (2.6) and (2.7) we get

$$\mathcal{L}_n\{\overline{\delta}_x^3 f(x); y\} = n^3 y^{3n} \mathcal{L}_n\{f(x); y\} - n^2 y^{2n} f(0^+) - ny^n \overline{\delta}_x f(0^+) - \overline{\delta}_x^2 f(0^+). \tag{2.8}$$

By repeated application of (2.6) and (2.8) we obtain the following theorem.

Theorem 2.1. If $f, f', ..., f^{(k-1)}$ are all continuous functions with a piecewise continuous derivative $f^{(k)}$ on the interval $[0,\infty)$, and if all functions are of exponential order $\exp(\alpha^n x^n)$ as $x \to \infty$ for some constant α then

$$\mathcal{L}_n\{\overline{\delta}_x^k f(x); y\} = (ny^n)^k \mathcal{L}_n\{f(x); y\} - (ny^n)^{k-1} f(0^+) - (ny^n)^{k-2}\overline{\delta}_x f(0^+) - \tag{2.9}$$

$$... - ny^n \overline{\delta}_x^{k-2} f(0^+) - \overline{\delta}_x^{k-1} f(0^+) \tag{2.10}$$

for $k \geq 1$, k is a positive integer.

The $\mathcal{L}_n$-transform defined in (1.3) is an analytic function in the half plane $Re\ y > \alpha$. Therefore, $\mathcal{L}_n\{f(x); y\}$ has derivatives of all orders and the derivatives can be formally obtained by differentiating (1.3). Applying the $\overline{\delta}$ with respect to the variable y we obtain

$$\overline{\delta}_y \mathcal{L}_n\{f(x); y\} = \frac{1}{y^{n-1}} \frac{d}{dy} \int_0^\infty x^{n-1}\exp(-y^n x^n) f(x) dx$$

$$= \frac{1}{y^{n-1}} \int_0^\infty x^{n-1}(-x^n n y^{n-1} \exp(-y^n x^n)) f(x) dx = -n\mathcal{L}_n\{x^n f(x); y\}. \tag{2.11}$$

If we keep taking the $\overline{\delta}$-derivative of (1.3) with respect to the variable y, then we deduce

$$\overline{\delta}_y^k \mathcal{L}_n\{f(x); y\} = \int_0^\infty x^{n-1} \overline{\delta}_y^k \exp(-y^n x^n) f(x) dx \tag{2.12}$$

for $k \in \mathbb{N}$.

$$\int_0^\infty x^{n-1} \overline{\delta}_y^k \exp(-y^n x^n) f(x) dx = \int_0^\infty x^{n-1} \overline{\delta}_y^{k-1} [(-n) x^n \exp(-y^n x^n)] f(x) dx$$

$$= \int_0^\infty x^{n-1} \overline{\delta}_y^{k-2} [(-n)^2 x^{2n} \exp(-y^n x^n)] f(x) dx$$

$$\cdots$$

$$= \int_0^\infty x^{n-1} [(-n)^k x^{kn} \exp(-y^n x^n)] f(x) dx = (-n)^k \mathcal{L}_n\{x^{kn} f(x); y\}. \tag{2.13}$$

As a result we obtain the following theorem.

Theorem 2.2. If f is piecewise continuous on the interval $[0, \infty)$ and is of exponential order $\exp(\alpha^n x^n)$ as $x \to \infty$, then

$$\mathcal{L}_n\{x^{kn} f(x); y\} = \frac{(-1)^k}{(n)^k} \overline{\delta}_y^k \mathcal{L}_n\{f(x); y\} \tag{2.14}$$

for $k \geq 1$, k is a positive integer.

Theorem 2.3. Let $\mathcal{L}_n\{f(x); y^{1/n}\}$, $n = 2^k$, $k = 0, 1, 2, ...$ be an analytic function of y except at singular points each of which lies to the left of the vertical line $Re\, y = a$ and they are finite numbers. Suppose that $y = 0$ is not a branch point and $\lim_{y \to \infty} \mathcal{L}_n\{f(x); y^{1/n}\} = 0$ in the left plane $Re\, y \leq a$ then, the following identity

$$\mathcal{L}_n^{-1}\{\mathcal{L}_n\{f(x); y\}\} = \frac{1}{2\pi i} \int_{a-i\infty}^{a+i\infty} n\mathcal{L}_n\{f(x); y^{1/n}\} \exp(yx^n) dy$$

$$= \sum_{k=1}^{m} [Res\{n\mathcal{L}_n\{f(x); y^{1/n}\} \exp(yx^n); y = y_k\}] \tag{2.15}$$

hold true for m singular points.

Proof. We take a vertical closed semi-circle as contour of integration. Using residues theorem and boundedness of $\mathcal{L}_n\{f(x); y^{1/n}\}$, we show that the identity (2.15) of Theorem 2.3 is valid. When $y = 0$ is a branch point we take key-hole contour instead of simple vertical semi-circle.

We assume that $\mathcal{L}_n\{f(x), y^{1/n}\}$ has a finite number of singularities in the left half plane $Rey \leqslant a$. Let $\gamma = \gamma_1 + \gamma_2$ be the closed contour consisting of the vertical line segment γ_1, which is defined from $a - iR$ to $a + iR$ and vertical semi-circle γ_2, that is defined as $|y - a| = R$. Let γ_2 lie to the left of vertical line γ_1. The radius R can be taken large enough so that γ encloses all the singularities of the $\mathcal{L}_n\{f(x); y^{1/n}\}$. Hence, by the residues theorem we have

$$\frac{1}{2\pi i}\int\limits_{a-i\infty}^{a+i\infty} n\mathcal{L}_n\{f(x); y^{1/n}\}\exp(yx^n)dy$$

$$= \frac{1}{2\pi i}\int\limits_{\gamma_1} n\mathcal{L}_n\{f(x); y^{1/n}\}\exp(yx^n)dy - \frac{1}{2\pi i}\int\limits_{\gamma_2} n\mathcal{L}_n\{f(x); y^{1/n}\}\exp(yx^n)dy$$

$$= \sum_{k=1}^{m}[Res\{n\mathcal{L}_n\{f(x); y^{1/n}\}\exp(yx^n); y = y_k\}] - \frac{1}{2\pi i}\int\limits_{\gamma_2} n\mathcal{L}_n\{f(x); y^{1/n}\}\exp(yx^n)dy \tag{2.16}$$

where $y_1, y_2, \ldots, y_m$ are all the singularities of $\mathcal{L}_n\{f(x); y^{1/n}\}$. Taking the limit from both sides of the relation (2.16) as R tends to $+\infty$, because of the Jordan's Lemma, the second integral in the right tends to zero.

Even $\mathcal{L}_n\{f(x); y^{1/n}\}$ has one branch point at $y = 0$, we can use the identity (2.15). The proof of the proposition is similar to the proof of the Main Theorem in the paper [1], where we take $n = 2^k, k \in \mathbb{N}$ instead of $n = 2$.

If the number of singularities is infinite, we take the semi-circles γ_m which is centered at point a, with radius $R_m = \pi^2 m^2, m \in \mathbb{N}$.

We illustrate the above Theorem by showing that the following examples.

Example 2.11. We show that

$$\mathcal{L}_n^{-1}\{\frac{1}{y^{2n} + a^{2n}}; x\} = \frac{n}{a^n}\sin(a^n x^n) \tag{2.17}$$

where $Re\ a > 0$.

Using the assertion (2.15) of Theorem 2.3 we obtain

$$\mathcal{L}_n^{-1}\{\frac{1}{y^{2n} + a^{2n}}; x\} = \sum_{k=1}^{2} Res[n\frac{1}{y^2 + a^{2n}}\exp(yx^n); y = y_k] \tag{2.18}$$

where the singular points are $y_k = \mp ia^n\ ,\ k = 1, 2$ and then we have

$$Res[n\frac{1}{y^2 + a^{2n}}\exp(yx^n); ia^n] = n\lim_{y\to ia^n}(y - ia^n)\frac{\exp(yx^n)}{y^2 + a^{2n}} = n\frac{\exp(ia^n x^n)}{2ia^n} \tag{2.19}$$

and similarly we have

$$Res[n\frac{1}{y^2+a^{2n}}\exp(yx^n);-ia^n]=-n\frac{\exp(-ia^nx^n)}{2ia^n}. \tag{2.20}$$

Using the relations (2.19) and (2.20) we find the formula (2.17) from (2.18) as follows

$$\mathcal{L}_n^{-1}\{\frac{1}{y^{2n}+a^{2n}};x\}=\frac{n}{a^n}\frac{\exp(ia^nx^n)-\exp(-ia^nx^n)}{2i}$$

$$=\frac{n}{a^n}\sin(a^nx^n). \tag{2.21}$$

Example 2.12. We show that

$$\mathcal{L}_n^{-1}\{\frac{1}{y^n}\exp(-\frac{a^n}{y^n});x\}=nJ_0(2a^{n/2}x^{n/2}) \tag{2.22}$$

where J_0 is the Bessel function of order zero.

Using the assertion (2.15) of Theorem 2.3 we have

$$\mathcal{L}_n^{-1}\{\frac{1}{y^n}\exp(-\frac{a^n}{y^n});x\}=Res[n\frac{1}{y}\exp(-\frac{a^n}{y})\exp(yx^n),y=y_k]. \tag{2.23}$$

From the following Taylor expansions of the exponential functions in (2.23),

$$n\frac{1}{y}\exp(-\frac{a^n}{y})\exp(yx^n)=\frac{n}{y}\sum_{m=0}^{\infty}(-1)^m\frac{a^{mn}}{m!y^m}\sum_{k=0}^{\infty}\frac{y^kx^{nk}}{k!}$$

$$=\frac{n}{y}[1-\frac{a^n}{1!y}+\frac{a^{2n}}{2!y^2}-\frac{a^{3n}}{3!y^3}+...][1+\frac{x^ny}{1!}+\frac{x^{2n}y^2}{2!}+\frac{x^{3n}}{3!}+...] \tag{2.24}$$

we find that $Res[n\frac{1}{y}\exp(\frac{a^n}{y})\exp(yx^n)]$ as the coefficient of the term $\frac{1}{y}$ as follows

$$Res[n\frac{1}{y}\exp(\frac{a^n}{y})\exp(yx^n)=n[1-\frac{a^nx^n}{(1!)^2}+\frac{a^{2n}x^{2n}}{(2!)^2}-\frac{a^{3n}x^{3n}}{(3!)^2}+...]$$

$$=n\sum_{m=0}^{\infty}(-1)^m\frac{(ax)^{mn}}{(n!)^2}=nJ_0(2a^{n/2}x^{n/2}). \tag{2.25}$$

Thus, we obtain from (2.25) and the formula (2.23), the assertion (2.22) of Example 2.12.

3. Application of the $\mathcal{L}_n$-transform to ordinary differential equations

First we consider the ordinary differential equation (1.53) for $v>n$ and $v=2^m+1\ ,m=0,1,2,...$

Dividing (1.53) by x^{n-1}, adding and subtracting the term $\frac{n-1}{x^{n-1}}z'$ we obtain

$$x^n\Big(\frac{1}{x^{2n-2}}z''-\frac{n-1}{x^{2n-1}}z'\Big)+\frac{n-1}{x^{n-1}}z'-\frac{2v+n-3}{x^{n-1}}z'+z=0. \tag{3.1}$$

Using the definition of the $\overline{\delta}$-derivative given in (2.1) and (2.2), we can express (3.1) as

$$x^n\overline{\delta}_x^2 z(x) - 2(v-1)\overline{\delta}_x z(x) + z(x) = 0. \tag{3.2}$$

Applying the $\mathcal{L}_n$-transform to (3.2) we find

$$\mathcal{L}_n\{x^n\overline{\delta}_x^2 z; y\} - 2(v-1)\mathcal{L}_n\{\overline{\delta}_x z; y\} + \mathcal{L}_n\{z(x); y\} = 0. \tag{3.3}$$

Using Theorem 2.1 for $k = 1$ and $k = 2$ in (3.3) and performing necessary calculations we obtain

$$-\frac{1}{n}\overline{\delta}_y\mathcal{L}_n\{\overline{\delta}_x^2 z; y\} - 2(v-1)\mathcal{L}_n\{\overline{\delta}_x z; y\} + \mathcal{L}_n\{z; y\} = 0, \tag{3.4}$$

$$-\frac{1}{n}\frac{1}{y^{n-1}}\frac{d}{dy}(n^2y^{2n}\overline{z}(y) - ny^n z(0^+) - \overline{\delta}_x z(0^+)) - 2(v-1)(ny^n\overline{z}(y) - z(0^+)) + \overline{z}(y) = 0 \tag{3.5}$$

where $\overline{z}(y) = \mathcal{L}_n\{z(x); y\}$. We assume that $z(0^+) = 0$. Thus, we obtain the following first order differential equation:

$$\overline{z}'(y) + (2(n+v-1)\frac{1}{y} - \frac{1}{ny^{n+1}})\overline{z}(y) = 0. \tag{3.6}$$

Solving the first order differential equation (3.6) we have

$$\overline{z}(y) = C\sum_{m=0}^{\infty}(-1)^m\frac{1}{m!n^{2m}y^{mn+2n+2v-2}}. \tag{3.7}$$

Applying the $\mathcal{L}_n^{-1}$ transform we obtain

$$z(x) = C\sum_{m=0}^{\infty}(-1)^m\frac{x^{mn+n+2v-2}}{m!\Gamma(m+\frac{n+2v-2}{n}+1)n^{2m-1}} \tag{3.8}$$

where we use the following relations

$$\mathcal{L}_n\{x^k; y\} = \frac{\Gamma(\frac{k}{n}+1)}{ny^{n+k}}\ ,\ k = mn+n+2v-2 \tag{3.9}$$

and

$$\mathcal{L}_n^{-1}\{\frac{1}{y^{mn+n+2v-2+n}}\} = \frac{nx^{mn+n+2v-2}}{\Gamma(m+1+\frac{2v-2}{n}+1)}. \tag{3.10}$$

Setting $\alpha = \frac{2v+n-2}{n}$, $C = n^{-\frac{2v-2}{n}-2}$ we obtain the solution of the ordinary differential equation (1.53)

$$z(x) = x^{\frac{n\alpha}{2}}J_\alpha(\frac{2}{n}x^{\frac{n}{2}}) \tag{3.11}$$

where $\alpha \in \mathbb{Z}$ because of the inequality $v > n$ $(v, n \in \mathbb{N})$ and J_α is the Bessel function of the first kind of order α.

In the second step we will use the $\mathcal{L}_n$-transform for solving (1.54).

Dividing (1.54) by x^{n-1}, adding and subtracting the term $\frac{n-1}{x^{n-1}}z'$ we obtain

$$x^n\Big(\frac{1}{x^{2n-2}z''(x) - \frac{n-1}{x^{2n-1}}z'(x)}\Big) + \frac{n-1}{x^{n-1}}z'(x) - (n^2-1)\frac{1}{x^{n-1}}z'(x) + z(x) = 0. \tag{3.12}$$

Using the definition of the $\overline{\delta}_x$-derivative (2.1) and (2.2) we can express (3.12) as

$$x^n\overline{\delta}_x^2 z(x) - n(n-1)\overline{\delta}_x z(x) + z(x) = 0. \tag{3.13}$$

Considering the following relations

$$\mathcal{L}_n\{x^n\overline{\delta}_x^2 z(x); y\} = -\frac{1}{n}\overline{\delta}_y\mathcal{L}_n\{\overline{\delta}_x^2 z(x); y\} = -2n^2y^n\overline{z}(y) - ny^{n+1}\overline{z}'(y) + nz(0^+), \tag{3.14}$$

$$n(n-1)\mathcal{L}_n\{\overline{\delta}_x z(x); y\} = n(n-1)(ny^n\overline{z}(y) - z(0^+)) = n^2(n-1)y^n\overline{z}(y) - n(n-1)z(0^+) \tag{3.15}$$

and applying the $\mathcal{L}_n$-transform to (3.13) we obtain

$$\mathcal{L}_n\{x^n\overline{\delta}_x^2 z(x); y\} - n(n-1)\mathcal{L}_n\{\overline{\delta}_x z(x); y\} + \mathcal{L}_n\{z(x); y\} = 0 \tag{3.16}$$

$$ny^{n+1}\overline{z}'(y) + [n^2(n+1)y^n - 1]\overline{z}(y) - n^2 z(0^+) = 0 \tag{3.17}$$

where $\overline{z}(y) = \mathcal{L}_n\{z(x); y\}$.

We may assume

$$z(0^+) = 0. \tag{3.18}$$

Solving the first order differential equation after substituting (3.18) into (3.17) we get

$$\overline{z}(y) = Cy^{-n^2-n}\exp(-\frac{1}{n^2y^n}). \tag{3.19}$$

Calculating the Taylor expansion of the exponential function in (3.19) we have

$$\overline{z}(y) = C\sum_{m=0}^{\infty}\frac{(-1)^m}{m!n^{2m}}\frac{1}{y^{n+nm+n^2}}. \tag{3.20}$$

Using the following relation

$$\mathcal{L}_n^{-1}\{\frac{1}{y^{n+nm+n^2}}\} = \frac{nx^{nm+n^2}}{\Gamma(m+n+1)} \tag{3.21}$$

and applying the $\mathcal{L}_n^{-1}$ transform to (3.20) we find

$$z(x) = Cn^{n+1}x^{\frac{n^2}{2}}\sum_{m=0}^{\infty}(-1)^m\frac{1}{m!\Gamma(m+n+1)}\Big(\frac{2x^{n/2}}{2n}\Big)^{2m+n}. \tag{3.22}$$

Setting $C = n^{-n-1}$ in (3.22) we obtain the solution of the equation (1.54)

$$z(x) = x^{\frac{n^2}{2}}J_n\{\frac{2}{n}x^{\frac{n}{2}}\} \tag{3.23}$$

where J_n is the Bessel function of the first kind of order n.

References

[1] Aghili A, Ansari A, Sedghi A.: An inversion technique for the $\mathcal{L}_2$-transform with applications. Int. J. Contemp. Math. Sciences 2.28, 1387-1394 (2007).

[2] Erdelyi, A.: Tables of lntegral Transforms vol. 1. New York,NY,USA, McGraw-Hill, (1954).

[3] Erdelyi, A.: Tables of lntegral Transforms vol. 2. New York,NY,USA, McGraw-Hill, (1954).

[4] Yürekli O, Sadek I.: A Parseval-Goldstein type theorem on the Widder potential transform and its applications. International Journal of Mathematics and Mathematical Sciences 14.3, 517-524 (1991).

[5] Yürekli O.: Theorems on $\mathcal{L}_2$-transform and its applications. Complex Variables and Elliptic Equations 38.2, 95-107 (1999).

[6] Yürekli O.: New identities involving the Laplace and the $\mathcal{L}_2$-transforms and their applications. Applied Mathematics and Computation 99.2, 141-151 (1999).

[7] Yürekli O, Wilson S.: A new method of solving Bessel's differential equation using the $\mathcal{L}_2$-transform. Applied Mathematics and Computation 130.2, 587-591 (2002).

[8] Yürekli O, Wilson S.: A new method of solving Hermite's differential equation using the $\mathcal{L}_2$-transform. Applied Mathematics and Computation 145.2, 495-500 (2003).